

anti- ABHD2 antibody

Product Information

Catalog No.: FNab00049

Size: 100μg Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

Clone ID: None IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

Background

Progesterone-dependent acylglycerol lipase that catalyzes hydrolysis of endocannabinoid arachidonoylglycerol(AG) from cell membrane(PubMed:26989199). Acts as a progesterone receptor: progesterone-binding activates the acylglycerol lipase activity, mediating degradation of 1-arachidonoylglycerol(1AG) and 2-arachidonoylglycerol(2AG) to glycerol and arachidonic acid(AA)(PubMed:26989199). Plays a key role in sperm capacitation in response to progesterone by mediating degradation of 2AG, an inhibitor of the sperm calcium channel CatSper, leading to calcium influx via CatSper and sperm activation(PubMed:26989199). May also play a role in smooth muscle cells migration(By similarity).

Immunogen information

Immunogen: abhydrolase domain containing 2

Synonyms: ABHD2, Protein PHPS1 2, HS1 2, Lung alpha/beta hydrolase 2, LABH2

Observed MW: 48 kDa UniprotID: P08910

Application

1

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)

Tel:(0086)027-87384275 Fax: (0086)027-87800889 <u>www.fn-test.com</u>




Reactivity: Human, Mouse, Rat


Tested Application: ELISA, WB, IHC

Recommended dilution: WB: 1:500-1:5000; IHC: 1:20-1:200

Image:



Immunohistochemistry of paraffin-embedded human spleen using FNab00049(ABHD2 antibody) at dilution of 1:100



HEK-293 cells were subjected to SDS PAGE followed by western blot with FNab00049(ABHD2 antibody) at dilution of 1:500

Wuhan Fine Biotech Co., Ltd.