

anti- AFF4 antibody

Product Information

Catalog No.: FNab00197

Size: 100µg Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

Clone ID: None IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

Background

Key component of the super elongation complex(SEC), a complex required to increase the catalytic rate of RNA polymerase II transcription by suppressing transient pausing by the polymerase at multiple sites along the DNA. In the SEC complex, AFF4 acts as a central scaffold that recruits other factors through direct interactions with ELL proteins(ELL, ELL2 or ELL3) and the P-TEFb complex. In case of infection by HIV-1 virus, the SEC complex is recruited by the viral Tat protein to stimulate viral gene expression.

Immunogen information

Immunogen: AF4/FMR2 family, member 4

Synonyms: AF4/FMR2 family member 4, AF4/FMR2 family, member 4, AF5Q31,

AFF4, MCEF, Protein AF 5q31

Observed MW: 127 kDa, observed 150 kDa

UniprotID: Q9UHB7

Application

Reactivity: Human, Mouse, Rat

1

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)


Tel:(0086)027-87384275 Fax: (0086)027-87800889 <u>www.fn-test.com</u>




Tested Application: ELISA, WB, IP

Recommended dilution: WB: 1:200-1:2000; IP: 1:200-1:1000

Image:



IP Result of anti-AFF4 (IP:FNab00197, 4ug; Detection:FNab00197 1:300) with HepG2 cells lysate 3200ug.



L02 cells were subjected to SDS PAGE followed by western blot with FNab00197(AFF4 antibody) at dilution of 1:500

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve. East Lake High-Tech Development Zone. Wuhan, Hubei, China (430206)

Tel:(0086)027-87384275 Fax: (0086)027-87800889 www.fn-test.com