

anti- ALKBH7 antibody

Product Information

Catalog No.: FNab00315

Size: 100μg Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

Clone ID: None IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

Background

May function as protein hydroxylase; can catalyze auto-hydroxylation at Leu-110(in vitro), but this activity may be due to the absence of the true substrate(PubMed:25122757). Required to induce programmed necrosis in response to DNA damage caused by cytotoxic alkylating agents. Acts by triggering the collapse of mitochondrial membrane potential and loss of mitochondrial function that leads to energy depletion and cell death(PubMed:23666923). ALKBH7-mediated necrosis is probably required to prevent the accumulation of cells with DNA damage(PubMed:23666923). Does not display DNA demethylase activity(PubMed:23666923). Involved in fatty acid metabolism(By similarity).

Immunogen information

Immunogen: alkB, alkylation repair homolog 7(E. coli)
Synonyms: ABH7, ALKBH7, SPATA11, UNQ6002

Observed MW: 25 kDa UniprotID: Q9BT30

Application

1

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)

Tel:(0086)027-87384275 Fax: (0086)027-87800889 <u>www.fn-test.com</u>




Reactivity: Human, Mouse, Rat


Tested Application: ELISA, WB, IHC

Recommended dilution: WB: 1:200-1:2000; IHC: 1:20-1:200

Image:



Immunohistochemistry of paraffin-embedded human testis tissue slide using FNab00315(ALKBH7 Antibody) at dilution of 1:50



human testis tissue were subjected to SDS PAGE followed by western blot with FNab00315(ALKBH7 antibody) at dilution of 1:500