

anti- BBS2 antibody

Product Information

Catalog No.: FNab00816

Size: 100μg Form: liquid

Purification: Protein A+G purification

Purity: ≥95% as determined by SDS-PAGE

Host: Mouse

Clonality: monoclonal

Clone ID: 9F10 IsoType: IgG2a

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

Background

The BBSome complex is thought to function as a coat complex required for sorting of specific membrane proteins to the primary cilia. The BBSome complex is required for ciliogenesis but is dispensable for centriolar satellite function. This ciliogenic function is mediated in part by the Rab8 GDP/GTP exchange factor, which localizes to the basal body and contacts the BBSome. Rab8(GTP) enters the primary cilium and promotes extension of the ciliary membrane. Firstly the BBSome associates with the ciliary membrane and binds to RAB3IP/Rabin8, the guanosyl exchange factor(GEF) for Rab8 and then the Rab8-GTP localizes to the cilium and promotes docking and fusion of carrier vesicles to the base of the ciliary membrane. The BBSome complex, together with the LTZL1, controls SMO ciliary trafficking and contributes to the sonic hedgehog(SHH) pathway regulation. Required for proper BBSome complex assembly and its ciliary localization.

Immunogen information

Immunogen: Bardet-Biedl syndrome 2

Synonyms: None
Observed MW: 80 kDa
Uniprot ID: Q9BXC9

1

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 Gaoxin Ave. East Lake High-Tech Development Zone. Wuhan, Hubei, China (430206)

Tel:(0086)027-87384275 Fax: (0086)027-87800889 <u>www.fn-test.com</u>




Application

Reactivity: Human, Dog
Tested Application: ELISA, WB, IF

Recommended dilution: WB: 1:500-1:2000; IF: 1:20-1:200

Image:



K-562 cells were subjected to SDS PAGE followed by western blot with FNab00816(BBS2 Antibody) at dilution of 1:1000