

# anti- BTK antibody

#### **Product Information**

Catalog No.: FNab00980

Size: 100μg Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

Clone ID: None IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

## **Background**

The protein encoded by this gene plays a crucial role in B-cell development. Mutations in this gene cause X-linked agammaglobulinemia type 1, which is an immunodeficiency characterized by the failure to produce mature B lymphocytes, and associated with a failure of Ig heavy chain rearrangement. Alternative splicing results in multiple transcript variants encoding different isoforms.

#### **Immunogen information**

Immunogen: Bruton agammaglobulinemia tyrosine kinase

Synonyms: AGMX1, ATK, BPK

Observed MW: 70-76 kDa
Uniprot ID: Q06187

### **Application**

Reactivity: Human, Mouse, Rat
Tested Application: ELISA, WB, IHC, IF

Recommended dilution: WB: 1:500 - 1:2000; IHC: 1:50 - 1:100; IF: 1:50 - 1:100

### Wuhan Fine Biotech Co., Ltd.


B9 Bld, High-Tech Medical Devices Park, No. 818 Gaoxin Ave. East Lake High-Tech Development Zone. Wuhan, Hubei, China (430206)

Tel:(0086)027-87384275 Fax: (0086)027-87800889 <u>www.fn-test.com</u>


1




# Image:



Immunohistochemistry of paraffin-embedded rat spleen using FNab00980(BTK antibody) at dilution of 1:100



Immunofluorescence analysis of HeLa cells using FNab00980(BTK antibody) at dilution of 1:50



mouse spleen tissue were subjected to SDS PAGE followed by western blot with FNab00980(BTK antibody) at dilution of 1:300

## Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 Gaoxin Ave. East Lake High-Tech Development Zone. Wuhan, Hubei, China (430206)

Tel:(0086)027-87384275 Fax: (0086)027-87800889 <u>www.fn-test.com</u>