

anti- CSDE1 antibody

Product Information

Catalog No.: FNab02011

Size: 100μg Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

Clone ID: None IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

Background

RNA-binding protein. Required for internal initiation of translation of human rhinovirus RNA. May be involved in translationally coupled mRNA turnover. Implicated with other RNA-binding proteins in the cytoplasmic deadenylation/translational and decay interplay of the FOS mRNA mediated by the major coding-region determinant of instability(mCRD) domain.

Immunogen information

Immunogen: cold shock domain containing E1, RNA-binding

Synonyms: D1S155E, KIAA0885, NRU, UNR

Observed MW: 89kd UniprotID: 075534

Application

Reactivity: Human, Mouse

Tested Application: ELISA, WB, IHC, IF

Recommended dilution: WB: 1:500-1:2000; IHC: 1:20-1:200; IF: 1:20-1:200

Wuhan Fine Biotech Co., Ltd.


B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)

Tel:(0086)027-87384275 Fax: (0086)027-87800889 www.fn-test.com


1



Image:



Immunohistochemistry of paraffin-embedded human skin cancer using FNab02011(UNR CSDE1 antibody) at dilution of 1:50



HeLa cells were subjected to SDS PAGE followed by western blot with FNab02011(UNR; CSDE1 antibody) at dilution of 1:1000