

anti- CSK antibody

Product Information

Catalog No.: FNab02016

Size: 100μg Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

Clone ID: None IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

Background

Non-receptor tyrosine-protein kinase that plays an important role in the regulation of cell growth, differentiation, migration and immune response. Phosphorylates tyrosine residues located in the C-terminal tails of Src-family kinases(SFKs) including LCK, SRC, HCK, FYN, LYN or YES1. Upon tail phosphorylation, Src-family members engage in intramolecular interactions between the phosphotyrosine tail and the SH2 domain that result in an inactive conformation. To inhibit SFKs, CSK is recruited to the plasma membrane via binding to transmembrane proteins or adapter proteins located near the plasma membrane. Suppresses signaling by various surface receptors, including T-cell receptor(TCR) and B-cell receptor(BCR) by phosphorylating and maintaining inactive several positive effectors such as FYN or LCK.

Immunogen information

Immunogen: c-src tyrosine kinase

Synonyms: None
Observed MW: 50 kDa
UniprotID: P41240

Application

1

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)

Tel:(0086)027-87384275 Fax: (0086)027-87800889 www.fn-test.com




Reactivity: Human


Tested Application: ELISA, WB, IHC, IF, IP

Recommended dilution: WB: 1:500-1:2000; IP: 1:200-1:1000; IHC: 1:20-1:200; IF: 1:20-1:200

Image:




Immunohistochemistry of paraffin-embedded human lung using FNab02016(CSK antibody) at dilution of 1:200



IP Result of anti-CSK (IP:FNab02016, 4ug; Detection:FNab02016 1:800) with HL-60 cells lysate 1000ug.

Wuhan Fine Biotech Co., Ltd.





HL-60 cells were subjected to SDS PAGE followed by western blot with FNab02016(CSK antibody) at dilution of 1:800