

anti- CYP4F11 antibody

Product Information

Catalog No.: FNab02170

Size: 100μg Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

Clone ID: None IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

Background

Omega-hydroxylase that oxidizes a variety of structurally unrelated compounds, including fatty acids and xenobiotics. Plays a key role in vitamin K catabolism by mediating omega-hydroxylation of vitamin K1(phylloquinone), and menaquinone-4(MK-4), a form of vitamin K2. Hydroxylation of phylloquinone and MK-4 probably regulates blood coagulation(PubMed:24138531). Catalyzes omega-hydroxylation of 3-hydroxy fatty acids, such as 3-hydroxypalmitate, 3-hydroxyoleate, 3-hydroxyarachidonate, and 3-hydroxystearate(PubMed:18065749, PubMed:19932081). Oxidizes drugs such as erythromycin, benzphetamine, ethylmorphine, chlorpromazine and imipramine(PubMed:15364545).

Immunogen information

Immunogen: cytochrome P450, family 4, subfamily F, polypeptide 11

Synonyms: None Observed MW: 57kd

UniprotID: Q9HBI6

Application

1

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)

Tel :(0086)027-87384275 Fax: (0086)027-87800889 <u>www.fn-test.com</u>




Reactivity: Human


Tested Application: ELISA, WB, IHC

Recommended dilution: WB: 1:500-1:2000; IHC: 1:20-1:200

Image:



Immunohistochemistry of paraffin-embedded human liver cancer using FNab02170(CYP4F11 antibody) at dilution of 1:100



human liver tissue were subjected to SDS PAGE followed by western blot with FNab02170(CYP4F11 antibody) at dilution of 1:500