

anti- DDA1 antibody

Product Information

| | |
|---------------|-----------------------------------------------------------------------------------------------------------------|
| Catalog No.: | FNab02282 |
| Size: | 100µg |
| Form: | liquid |
| Purification: | Immunogen affinity purified |
| Purity: | ≥95% as determined by SDS-PAGE |
| Host: | Rabbit |
| Clonality: | polyclonal |
| Clone ID: | None |
| IsoType: | IgG |
| Storage: | PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months (Avoid repeated freeze / thaw cycles.) |

Background

May be involved in ubiquitination and subsequent proteasomal degradation of target proteins. Component of the DDD-E2 complexes which may provide a platform for interaction with CUL4A and WD repeat proteins.

Immunogen information

| | |
|--------------|----------------------------|
| Immunogen: | DET1 and DDB1 associated 1 |
| Synonyms: | C19orf58, PCIA1 |
| Observed MW: | 12 kDa |
| UniprotID : | Q9BW61 |

Application


| | |
|---------------------------|-------------------|
| Reactivity: | Human, Mouse, Rat |
| Tested Application: | ELISA, WB |
| Recommended dilution: WB: | 1:500-1:2000 |
| Image: | |

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 Gaoxin Ave. East Lake High-Tech Development Zone. Wuhan, Hubei, China (430206)

Tel : (0086)027-87384275

Fax: (0086)027-87800889 www.fn-test.com



HepG2 cells were subjected to SDS PAGE followed by western blot with FNab02282(DDA1 antibody) at dilution of 1:100