

anti- DIS3L antibody

Product Information

Catalog No.: FNab02395

Size: 100μg Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

Clone ID: None IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

Background

Putative cytoplasm-specific catalytic component of the RNA exosome complex which has 3'->5' exoribonuclease activity and participates in a multitude of cellular RNA processing and degradation events. In the cytoplasm, the RNA exosome complex is involved in general mRNA turnover and specifically degrades inherently unstable mRNAs containing AU-rich elements(AREs) within their 3' untranslated regions, and in RNA surveillance pathways, preventing translation of aberrant mRNAs. It seems to be involved in degradation of histone mRNA.

Immunogen information

Immunogen: DIS3 mitotic control homolog(S. cerevisiae)-like

Synonyms: DIS3L1, KIAA1955

Observed MW: 121 kDa UniprotID: Q8TF46

Application

Reactivity: Human

1

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)


Tel:(0086)027-87384275 Fax: (0086)027-87800889 <u>www.fn-test.com</u>



Tested Application: ELISA, WB

Recommended dilution: WB: 1:500-1:2000

Image:



K-562 cells were subjected to SDS PAGE followed by western blot with FNab02395(DIS3L Antibody) at dilution of 1:100

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve. East Lake High-Tech Development Zone. Wuhan, Hubei, China (430206)

Tel: (0086)027-87384275 Fax: (0086)027-87800889 www.fn-test.com