

anti- EHD2 antibody

Product Information

Catalog No.: FNab02675

Size: 100μg Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

Clone ID: None IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

Background

ATP-and membrane-binding protein that controls membrane reorganization/tubulation upon ATP hydrolysis(By similarity). Plays a role in membrane trafficking between the plasma membrane and endosomes(PubMed:17233914). Important for the internalization of GLUT4.Required for fusion of myoblasts to skeletal muscle myotubes. Required for normal translocation of FER1L5 to the plasma membrane(By similarity).

Immunogen information

Immunogen: EH-domain containing 2

Synonyms: PAST2
Observed MW: 61 kDa
UniprotID: O9NZN4

Application

Reactivity: Human, Mouse, Rat
Tested Application: ELISA, WB, IHC, IP

Recommended dilution: WB: 1:500-1:2000; IP: 1:200-1:1000; IHC: 1:20-1:200

Wuhan Fine Biotech Co., Ltd.


B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)

Tel:(0086)027-87384275 Fax: (0086)027-87800889 www.fn-test.com


1



Image:




Immunohistochemistry of paraffin-embedded human lung cancer using FNab02675(EHD2 antibody) at dilution of 1:50



IP Result of anti-EHD2 (IP:FNab02675, 4ug; Detection:FNab02675 1:800) with mouse heart tissue lysate 4000ug.





human heart tissue were subjected to SDS PAGE followed by western blot with FNab02675(EHD2 antibody) at dilution of 1:400

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve. East Lake High-Tech Development Zone. Wuhan, Hubei, China (430206)

Tel: (0086)027-87384275 Fax: (0086)027-87800889 www.fn-test.com