

anti- EHD3 antibody

Product Information

Catalog No.:	FNab02676
Size:	100µg
Form:	liquid
Purification:	Immunogen affinity purified
Purity:	≥95% as determined by SDS-PAGE
Host:	Rabbit
Clonality:	polyclonal
Clone ID:	None
IsoType:	IgG
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months (Avoid repeated freeze / thaw cycles.)

Background

ATP-and membrane-binding protein that controls membrane reorganization/tubulation upon ATP hydrolysis(PubMed:25686250). In vitro causes tubulation of endocytic membranes(PubMed:24019528). Binding to phosphatidic acid induces its membrane tubulation activity(By similarity). Plays a role in endocytic transport. Involved in early endosome to recycling endosome compartment(ERC), retrograde early endosome to Golgi, and endosome to plasma membrane(rapid recycling) protein transport. Involved in the regulation of Golgi maintenance and morphology(PubMed:16251358, PubMed:17233914, PubMed:19139087, PubMed:23781025). Involved in the recycling of internalized D1 dopamine receptor(PubMed:21791287). Plays a role in cardiac protein trafficking probably implicating ANK2(PubMed:20489164). Involved in the ventricular membrane targeting of SLC8A1 and CACNA1C and probably the atrial membrane localization of CACNA1GG and CACNA1H implicated in the regulation of atrial myocyte excitability and cardiac conduction(By similarity). In conjunction with EHD4 may be involved in endocytic trafficking of KDR/VEGFR2 implicated in control of glomerular function(By similarity). Involved in the rapid recycling of integrin beta-3 implicated in cell adhesion maintenance(PubMed:23781025). Involved in the unidirectional retrograde dendritic transport of endocytosed BACE1 and in efficient sorting of BACE1 to axons implicating a function in neuronal APP processing(By similarity). Plays a role in the formation of the ciliary vesicle, an early step in cilium biogenesis; possibly sharing redundant functions with EHD1(PubMed:25686250).

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 Gaoxin Ave. East Lake High-Tech Development Zone. Wuhan, Hubei, China(430206)

Tel : (0086)027-87384275


Fax: (0086)027-87800889 www.fn-test.com

Immunogen information

Immunogen: EH-domain containing 3
Synonyms: EHD2,
Observed MW: 61 kDa
UniprotID : Q9NZN3

Application

Reactivity: Human, Mouse
Tested Application: ELISA, WB
Recommended dilution: WB: 1:500-1:2000
Image:



mouse brain tissue were subjected to SDS PAGE followed by western blot with FNab02676(EHD3 Antibody) at dilution of 1:300

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve. East Lake High-Tech Development Zone. Wuhan, Hubei, China(430206)

Tel : (0086)027-87384275

Fax: (0086)027-87800889 www.fn-test.com