

anti- EIF3J antibody

Product Information

Catalog No.:	FNab02711
Size:	100µg
Form:	liquid
Purification:	Immunogen affinity purified
Purity:	\geq 95% as determined by SDS-PAGE
Host:	Rabbit
Clonality:	polyclonal
Clone ID:	None
IsoType:	IgG
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months (Avoid repeated freeze / thaw cycles.)

Background

Component of the eukaryotic translation initiation factor 3(eIF-3) complex, which is required for several steps in the initiation of protein synthesis. The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl-tRNAi and eIF-5 to form the 43S preinitiation complex(43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of post-termination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation. This subunit binds directly within the mRNA entry channel of the 40S ribosome to the aminoacyl(A) site. It may regulate the interaction between the 43S PIC and mRNA.

Immunogen information

Immunogen:	eukaryotic translation initiation factor 3, subunit J
Synonyms:	EIF3S1
Observed MW:	35 kDa
UniprotID :	075822

Application


Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)

Tel :(0086)027-87384275 Fax: (0086)027-87800889 <u>www.fn-test.com</u>



Reactivity:Human, Mouse, RatTested Application:ELISA, WB, IHC, IF, IPRecommended dilution:WB: 1:500-1:2000; IP: 1:200-1:1000; IHC: 1:20-1:200; IF: 1:20-1:200Image:



Immunohistochemistry of paraffin-embedded human breast cancer using FNab02711(EIF3J antibody) at dilution of 1:100


IP Result of anti-EIF3J (IP:FNab02711, 4ug; Detection:FNab02711 1:500) with HeLa cells lysate 880ug.

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)

Tel :(0086)027-87384275 Fax: (0086)027-87800889 www.fn-test.com





HeLa cells were subjected to SDS PAGE followed by western blot with FNab02711(EIF3J antibody) at dilution of 1:500

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)

Tel :(0086)027-87384275 Fax: (0086)027-87800889 www.fn-test.com