

anti- EXD2 antibody

Product Information

Catalog No.: FNab02890

Size: 100μg Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

Clone ID: None IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

Background

Exonuclease required for double-strand breaks resection and efficient homologous recombination. Plays a key role in controlling the initial steps of chromosomal break repair, it is recruited to chromatin in a damage-dependent manner and functionally interacts with the MRN complex to accelerate resection through its 3'-5' exonuclease activity, which efficiently processes double-stranded DNA substrates containing nicks.

Immunogen information

Immunogen: exonuclease 3'-5' domain containing 2

Synonyms: C14orf114,

Observed MW: 56 kDa

UniprotID: Q9NVH0

Application

Reactivity: Human, Mouse, Rat

Tested Application: ELISA, WB, IF, IHC, IP

Recommended dilution: WB: 1:500-1:2000; IP: 1:200-1:1000; IHC: 1:20-1:200; IF: 1:20-1:200

Wuhan Fine Biotech Co., Ltd.


B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve. East Lake High-Tech Development Zone. Wuhan, Hubei, China (430206)

Tel :(0086)027-87384275 Fax: (0086)027-87800889 <u>www.fn-test.com</u>


1



Image:




Immunohistochemistry of paraffin-embedded human testis using FNab02890(EXD2 antibody) at dilution of 1:50



IP Result of anti-EXD2 (IP:FNab02890, 3ug; Detection:FNab02890 1:300) with mouse lung tissue lysate 5160ug.





mouse lung tissue were subjected to SDS PAGE followed by western blot with FNab02890(EXD2 antibody) at dilution of 1:300