

# anti- Histone H1.2 antibody

#### **Product Information**

Catalog No.: FNab03883

Size: 100μg Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

Clone ID: None IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

## **Background**

Histone H1 regulates chromatin competency through its high affinity binding to linker DNA as a structural component. It acts also as a regulator of individual gene transcription through chromatin remodeling, nucleosome spacing and DNA methylation. The calcualted molecular weight of Histone H1.2 is 21 kDa, but modified Histone H1.2 is about 32 kDa.

## **Immunogen information**

Immunogen: histone cluster 1, H1c

Synonyms: H1.2, H1F2, HIST1H1C, histone cluster 1, H1c, Histone H1.2, Histone

H<sub>1</sub>d

Observed MW: 32kd

UniprotID: P16403

### **Application**

Reactivity: Human, Mouse, Rat

Tested Application: ELISA, IHC, WB, IF, IP

Recommended dilution: WB: 1:500-1:3000; IP: 1:500-1:1000; IHC: 1:20-1:200; IF: 1:10-1:100

### Wuhan Fine Biotech Co., Ltd.


B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)

Tel:(0086)027-87384275 Fax: (0086)027-87800889 www.fn-test.com


1




## Image:



Immunohistochemical of paraffin-embedded human ovary tumor using FNab03883(HIST1H1C antibody) at dilution of 1:100



Immunofluorescent analysis of Hela cells, using HIST1H1C antibody FNab03883 at 1:25 dilution and Rhodamine-labeled goat anti-rabbit IgG (red).




IP Result of anti-Histone H1.2 (IP: FNab03883, 3ug; Detection: FNab03883 1:600) with HeLa cells lysate 1200ug.

## Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve. East Lake High-Tech Development Zone. Wuhan, Hubei, China (430206)

Tel: (0086)027-87384275 Fax: (0086)027-87800889 www.fn-test.com





L02 cells were subjected to SDS PAGE followed by western blot with FNab03883(HIST1H1C antibody) at dilution of 1:500