

anti- HNMT antibody

Product Information

Catalog No.: FNab03944

Size: 100μg Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

Clone ID: None IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

Background

In mammals, histamine is metabolized by two major pathways: N(tau)-methylation via histamine N-methyltransferase and oxidative deamination via diamine oxidase. This gene encodes the first enzyme which is found in the cytosol and uses S-adenosyl-L-methionine as the methyl donor. In the mammalian brain, the neurotransmitter activity of histamine is controlled by N(tau)-methylation as diamine oxidase is not found in the central nervous system. A common genetic polymorphism affects the activity levels of this gene product in red blood cells. Multiple alternatively spliced transcript variants that encode different proteins have been found for this gene.

Immunogen information

Immunogen: histamine N-methyltransferase

Synonyms: HMT
Observed MW: 27 kDa
Uniprot ID: P50135

Application

1

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 Gaoxin Ave. East Lake High-Tech Development Zone. Wuhan, Hubei, China (430206)

Tel :(0086)027-87384275 Fax: (0086)027-87800889 <u>www.fn-test.com</u>




Reactivity: Human, Mouse


Tested Application: ELISA, WB, IHC, IP

Recommended dilution: WB: 1:500 - 1:2000; IHC: 1:50 - 1:200; IP: 1:20 - 1:50

Image:



Immunohistochemistry of paraffin-embedded mouse heart using FNab03944(HNMT antibody) at dilution of 1:50



human liver tissue were subjected to SDS PAGE followed by western blot with FNab03944(HNMT antibody) at dilution of 1:1000

Wuhan Fine Biotech Co., Ltd.