

anti- HSD3B7 antibody

Product Information

Catalog No.: FNab04032

Size: $100\mu g$ Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

Clone ID: None IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

Background

The 3-beta-HSD enzymatic system plays a crucial role in the biosynthesis of all classes of hormonal steroids. HSD VII is active against four 7-alpha-hydroxylated sterols. Does not metabolize several different C(19/21) steroids as substrates. Involved in bile acid synthesis(PubMed:11067870). Plays a key role in cell positioning and movement in lymphoid tissues by mediating degradation of 7-alpha,25-dihydroxycholesterol(7-alpha,25-OHC): 7-alpha,25-OHC acts as a ligand for the G protein-coupled receptor GPR183/EBI2, a chemotactic receptor for a number of lymphoid cells.

Immunogen information

Immunogen: hydroxy-delta-5-steroid dehydrogenase, 3 beta-and steroid delta-

isomerase 7

Synonyms: None
Observed MW: 36 kDa
UniprotID: Q9H2F3

Application

1

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)

Tel:(0086)027-87384275 Fax: (0086)027-87800889 <u>www.fn-test.com</u>




Reactivity: Human, Mouse


Tested Application: ELISA, WB, IHC

Recommended dilution: WB: 1:500-1:2000; IHC: 1:20-1:200

Image:



Immunohistochemistry of paraffin-embedded human pancreas cancer using FNab04032(HSD3B7 antibody) at dilution of 1:50



mouse liver tissue were subjected to SDS PAGE followed by western blot with FNab04032(HSD3B7 antibody) at dilution of 1:500