

anti- IL17RC antibody

Product Information

Catalog No.:	FNab04231
Size:	100µg
Form:	liquid
Purification:	Immunogen affinity purified
Purity:	≥95% as determined by SDS-PAGE
Host:	Rabbit
Clonality:	polyclonal
Clone ID:	None
IsoType:	IgG
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months (Avoid repeated freeze / thaw cycles.)

Background

Receptor for IL17A and IL17F homodimers as part of a heterodimeric complex with IL17RA(PubMed:16785495). Receptor for the heterodimer formed by IL17A and IL17B as part of a heterodimeric complex with IL17RA(PubMed:18684971). Has also been shown to be the cognate receptor for IL17F and to bind IL17A with high affinity without the need for IL17RA(PubMed:17911633). Activation of IL17RC leads to induction of expression of inflammatory chemokines and cytokines such as CXCL1(PubMed:16785495). Isoform 5: Receptor for both IL17A and IL17F. Isoform 6: Does not bind IL17A or IL17F. Isoform 7: Does not bind IL17A or IL17F. Isoform 8: Receptor for both IL17A and IL17F.

Immunogen information

Immunogen:	interleukin 17 receptor C
Synonyms:	None
Observed MW:	45-50kd
UniprotID :	Q8NAC3

Application

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)

Tel : (0086)027-87384275


Fax: (0086)027-87800889 www.fn-test.com

Reactivity: Human, Mouse

Tested Application: ELISA, WB

Recommended dilution: WB: 1:500-1:2000

Image:



HEK-293 cells were subjected to SDS PAGE followed by western blot with 1FNab04231(IL17RC antibody) at dilution of 1:200

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve. East Lake High-Tech Development Zone. Wuhan, Hubei, China(430206)

Tel : (0086)027-87384275

Fax: (0086)027-87800889 www.fn-test.com