

anti- IST1 antibody

Product Information

Catalog No.:	FNab04408
Size:	100µg
Form:	liquid
Purification:	Immunogen affinity purified
Purity:	\geq 95% as determined by SDS-PAGE
Host:	Rabbit
Clonality:	polyclonal
Clone ID:	None
IsoType:	IgG
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months (Avoid repeated freeze / thaw cycles.)

Background

ESCRT-III-like protein involved in specific functions of the ESCRT machinery. Is required for efficient abscission during cytokinesis, but not for HIV-1 budding. The involvement in the MVB pathway is not established. Involved in recruiting VPS4A and/or VPS4B to the midbody of dividing cells(PubMed:19129479, PubMed:19129480). During late anaphase, involved in nuclear envelope reassembly and mitotic spindle disassembly together with the ESCRT-III complex: IST1 acts by mediating the recruitment of SPAST to the nuclear membrane, leading to microtubule severing(PubMed:26040712). Regulates early endosomaltubulation together with the ESCRT-III complex by mediating the recruitment of SPAST(PubMed:23897888).

Immunogen information

4
1

Application

Wuhan Fine Biotech Co., Ltd.


B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)


Tel :(0086)027-87384275 Fax: (0086)027-87800889 <u>www.fn-test.com</u>

1



Reactivity:HumanTested Application:ELISA, WB, IF, IHC, IPRecommended dilution:WB: 1:500-1:2000; IP: 1:200-1:1000; IHC: 1:20-1:200; IF: 1:20-1:200Image:





Immunohistochemistry of paraffin-embedded human lung cancer using FNab04408(hIST1 antibody) at dilution of 1:200

human skeletal muscle tissue were subjected to SDS PAGE followed by western blot with FNab04408(hIST1 antibody) at dilution of 1:300

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)

Tel :(0086)027-87384275 Fax: (0086)027-87800889 <u>www.fn-test.com</u>