

anti- LZTS2 antibody

Product Information

Catalog No.: FNab04918

Size: 100μg Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

Clone ID: None IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

Background

Negative regulator of katanin-mediated microtubule severing and release from the centrosome. Required for central spindle formation and the completion of cytokinesis. May negatively regulate axonal outgrowth by preventing the formation of microtubule bundles that are necessary for transport within the elongating axon. Negative regulator of the Wnt signaling pathway. Represses beta-catenin-mediated transcriptional activation by promoting the nuclear exclusion of beta-catenin.

Immunogen information

Immunogen: leucine zipper, putative tumor suppressor 2

Synonyms: KIAA1813, LAPSER1

Observed MW: 70-80 kDa UniprotID: Q9BRK4

Application

Reactivity: Human, Mouse, Rat Tested Application: ELISA, IHC, WB

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)


Tel:(0086)027-87384275 Fax: (0086)027-87800889 www.fn-test.com

1




Recommended dilution: WB: 1:500-1:2000; IHC: 1:20-1:200

Image:



Immunohistochemistry of paraffin-embedded human prostate hyperplasia tissue slide using FNab04918(LZTS2 Antibody) at dilution of 1:50



PC-3 cells were subjected to SDS PAGE followed by western blot with FNab04918(LZTS2 Antibody) at dilution of 1:300