

anti- MAFF antibody

Product Information

Catalog No.: FNab04931

Size: 100μg Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

Clone ID: None IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

Background

Interacts with the upstream promoter region of the oxytocin receptor gene. May be a transcriptional enhancer in the up-regulation of the oxytocin receptor gene at parturition. Since it lacks a putative transactivation domain, it may behave as a transcriptional repressor when it dimerize among himself. May also serve as a transcriptional activator by dimerizing with other(usually larger) basic-zipper proteins and recruiting them to specific DNA-binding sites. May be involved in the cellular stress response.

Immunogen information

Immunogen: v-mafmusculoaponeuroticfibrosarcoma oncogene homolog F(avian)

Synonyms: None

Observed MW: 18-23 kDa UniprotID: Q9ULX9

Application

Reactivity: Human, Mouse, Rat Tested Application: ELISA, WB, IHC

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)


Tel:(0086)027-87384275 Fax: (0086)027-87800889 www.fn-test.com

1




Recommended dilution: WB: 1:500-1:2000; IHC: 1:20-1:200

Image:



Immunohistochemistry of paraffin-embedded human ovary tumor using FNab04931(MAFF antibody) at dilution of 1:100



human lung tissue were subjected to SDS PAGE followed by western blot with FNab04931(MAFF antibody) at dilution of 1:300