

# anti- MED10 antibody

# **Product Information**

Catalog No.:	FNab05086
Size:	100µg
Form:	liquid
Purification:	Immunogen affinity purified
Purity:	$\geq$ 95% as determined by SDS-PAGE
Host:	Rabbit
Clonality:	polyclonal
Clone ID:	None
IsoType:	IgG
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months (Avoid repeated freeze / thaw cycles.)

# Background

Component of the Mediator complex, a coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors.

### Immunogen information

Immunogen:	mediator complex subunit 10
Synonyms:	L6, MED10, mediator complex subunit 10, NUT2, TRG 17, TRG 20, TRG17, TRG20
Observed MW:	16 kDa
Uniprot ID :	Q9BTT4

# Application

Reactivity: Human, Mouse, Rat

### Wuhan Fine Biotech Co., Ltd.


B9 Bld, High-Tech Medical Devices Park, No. 818 Gaoxin Ave.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)

Tel :( 0086)027-87384275 Fax: (0086)027-87800889

www.fn-test.com



Tested Application: ELISA, WB Recommended dilution: WB: 1:200 - 1:1000 Image:



COLO320 cells were subjected to SDS PAGE followed by western blot with FNab05086( MED10 Antibody) at dilution of 1:500

#### Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 Gaoxin Ave.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)

www.fn-test.com