

anti- MYO16 antibody

Product Information

| Catalog No.: | FNab05499 |
|---------------|---|
| Size: | 100µg |
| Form: | liquid |
| Purification: | Immunogen affinity purified |
| Purity: | \geq 95% as determined by SDS-PAGE |
| Host: | Rabbit |
| Clonality: | polyclonal |
| Clone ID: | None |
| IsoType: | IgG |
| Storage: | PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months (Avoid repeated freeze / thaw cycles.) |

Background

Myosins are actin-based motor molecules with ATPase activity. Unconventional myosins serve in intracellular movements. Their highly divergent tails are presumed to bind to membranous compartments, which would be moved relative to actin filaments. May be involved in targeting of the catalytic subunit of protein phosphatase 1 during brain development. Activates PI3K and concomitantly recruits the WAVE1 complex to the close vicinity of PI3K and regulates neuronal morphogenesis(By similarity).

Immunogen information

| Immunogen: | myosin XVI |
|--------------|-------------|
| Synonyms: | KIAA0865, |
| Observed MW: | 200-240 kDa |
| UniprotID : | Q9Y6X6 |

Application

| Reactivity: | Human, Mouse |
|---------------------|----------------|
| Tested Application: | ELISA, WB, IHC |


Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)

Tel :(0086)027-87384275 Fax: (0086)027-87800889 www.fn-test.com



Recommended dilution: WB: 1:500-1:2000; IHC: 1:20-1:200 Image:



Immunohistochemistry of paraffin-embedded human bladder using FNab05499(MYO16 antibody) at dilution of 1:50

human brain tissue were subjected to SDS PAGE followed by western blot with FNab05499(MYO16 antibody) at dilution of 1:500

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)

Tel :(0086)027-87384275 Fax: (0086)027-87800889 <u>www.fn-test.com</u>