

anti- NKD2 antibody

Product Information

| Catalog No.: | FNab05743 |
|---------------|---|
| Size: | 100µg |
| Form: | liquid |
| Purification: | Immunogen affinity purified |
| Purity: | \geq 95% as determined by SDS-PAGE |
| Host: | Rabbit |
| Clonality: | polyclonal |
| Clone ID: | None |
| IsoType: | IgG |
| Storage: | PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months (Avoid repeated freeze / thaw cycles.) |

Background

Cell autonomous antagonist of the canonical Wnt signaling pathway. May activate a second Wnt signaling pathway that controls planar cell polarity(By similarity). Required for processing of TGFA and for targeting of TGFA to the basolateral membrane of polarized epithelial cells.

Immunogen information

| Immunogen: | naked cuticle homolog 2(Drosophila) |
|--------------|-------------------------------------|
| Synonyms: | None |
| Observed MW: | 59kd |
| UniprotID : | Q969F2 |

Application


| Reactivity: | Human, Mouse, Rat | | |
|--|-------------------|--|--|
| Tested Application: | ELISA, WB | | |
| Recommended dilution: WB: 1:500-1:2000 | | | |
| Image: | | | |

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)

Tel :(0086)027-87384275 Fax: (0086)027-87800889 www.fn-test.com





mouse kidney tissue were subjected to SDS PAGE followed by western blot with FNab05743(NKD2 antibody) at dilution of 1:400

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)

Tel :(0086)027-87384275 Fax: (0086)027-87800889 <u>www.fn-test.com</u>