

# anti- NSFL1C antibody

#### **Product Information**

Catalog No.: FNab05868

Size: 100μg Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

Clone ID: None IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

### **Background**

Reduces the ATPase activity of VCP. Necessary for the fragmentation of Golgi stacks during mitosis and for VCP-mediated reassembly of Golgi stacks after mitosis. May play a role in VCP-mediated formation of transitional endoplasmic reticulum(tER)(By similarity). Inhibits the activity of CTSL(in vitro).

### **Immunogen information**

Immunogen: NSFL1(p97) cofactor(p47)

Synonyms: UBXN2C

Observed MW: 47 kDa, 41 kDa

UniprotID: Q9UNZ2

#### **Application**

Reactivity: Human, Mouse Tested Application: ELISA, WB

Recommended dilution: WB: 1:500-1:2000

#### Wuhan Fine Biotech Co., Ltd.


B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)

Tel:(0086)027-87384275 Fax: (0086)027-87800889 www.fn-test.com

1



## Image:



A431 cells were subjected to SDS PAGE followed by western blot with FNab05868(NSFL1C antibody) at dilution of 1:800

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve. East Lake High-Tech Development Zone. Wuhan, Hubei, China (430206)

Tel: (0086)027-87384275 Fax: (0086)027-87800889 www.fn-test.com