

anti-Peroxiredoxin 3 antibody

Product Information

Catalog No.: FNab06322

Size: 100μg Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

Clone ID: None IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

Background

This gene encodes a mitochondrial protein with antioxidant function. The protein is similar to the C22 subunit of Salmonella typhimurium alkylhydroperoxide reductase, and it can rescue bacterial resistance to alkylhydroperoxide in E. coli that lack the C22 subunit. The human and mouse genes are highly conserved, and they map to the regions syntenic between mouse and human chromosomes. Sequence comparisons with recently cloned mammalian homologs suggest that these genes consist of a family that is responsible for the regulation of cellular proliferation, differentiation and antioxidant functions. This family member can protect cells from oxidative stress, and it can promote cell survival in prostate cancer. Alternative splicing of this gene results in multiple transcript variants. Related pseudogenes have been identified on chromosomes 1, 3, 13 and 22.

Immunogen information

Immunogen: peroxiredoxin 3

Synonyms: Antioxidant protein 1, AOP 1, AOP1, HBC189, MER5, Peroxiredoxin 3,

Peroxiredoxin III, PRDX3, PRO1748, Protein MER5 homolog, Prx III,

PRX3, SP 22

Observed MW: 25 kDa
UniprotID: P30048

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)

Tel:(0086)027-87384275 Fax: (0086)027-87800889 www.fn-test.com

1




Application


Reactivity: Human, Mouse, Rat Tested Application: ELISA, WB, IHC, IF

Recommended dilution: WB: 1:500 - 1:2000; IHC: 1:50 - 1:200; IF: 1:50 - 1:200

Image:




Immunohistochemistry of paraffin-embedded human liver cancer tissue slide using FNab06322(Prx III Antibody) at dilution of 1:50



Immunofluorescence analysis of HeLa cells using FNab06322(Prx III Antibody). Blue: DAPI for nuclear staining.





HeLa cells were subjected to SDS PAGE followed by western blot with FNab06322(Prx III antibody) at dilution of 1:1000

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve. East Lake High-Tech Development Zone. Wuhan, Hubei, China (430206)

Tel: (0086)027-87384275 Fax: (0086)027-87800889 www.fn-test.com