

# anti- PKC alpha antibody

#### **Product Information**

Catalog No.: FNab06480

Size: 100μg Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

## **Background**

Protein kinase C (PKC) is a family of serine- and threonine-specific protein kinases that can be activated by calcium and the second messenger diacylglycerol. PKC family members phosphorylate a wide variety of protein targets and are known to be involved in diverse cellular signaling pathways. PKC family members also serve as major receptors for phorbol esters, a class of tumor promoters. Each member of the PKC family has a specific expression profile and is believed to play a distinct role in cells. The protein encoded by this gene is one of the PKC family members. This kinase has been reported to play roles in many different cellular processes, such as cell adhesion, cell transformation, cell cycle checkpoint, and cell volume control. Knockout studies in mice suggest that this kinase may be a fundamental regulator of cardiac contractility and Ca(2+) handling in myocytes.

## **Immunogen information**

Immunogen: protein kinase C, alpha

Synonyms: AAG6, PKC A, PKC alpha, PKCA, PRKACA, PRKCA, Protein kinase

C alpha type, protein kinase C, alpha

Observed MW: 77kDa Uniprot ID: P17252

**Application** 

1

### Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 Gaoxin Ave. East Lake High-Tech Development Zone. Wuhan, Hubei, China (430206)

Tel :( 0086)027-87384275 Fax: (0086)027-87800889 <u>www.fn-test.com</u>




Reactivity: Human


Tested Application: ELISA, WB, IHC, IF

Recommended dilution: WB: 1:500 - 1:2000; IHC: 1:50 - 1:200; IF: 1:50 - 1:200


Image:



Immunohistochemistry of paraffin-embedded human thyroid cancer tissue slide using FNab06480( PRKCA Antibody) at dilution of 1:200



Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using FNab06480( PRKCA Antibody) at dilution of 1:50 and Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L)



Jurkat cells were subjected to SDS PAGE followed by western blot with FNab06480(PRKCA antibody) at dilution of 1:1000