

# anti- PRKACB antibody

#### **Product Information**

Catalog No.: FNab06778

Size: 100μg Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

Clone ID: None IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

### **Background**

Mediates cAMP-dependent signaling triggered by receptor binding to GPCRs. PKA activation regulates diverse cellular processes such as cell proliferation, the cell cycle, differentiation and regulation of microtubule dynamics, chromatin condensation and decondensation, nuclear envelope disassembly and reassembly, as well as regulation of intracellular transport mechanisms and ion flux. Regulates the abundance of compartmentalized pools of its regulatory subunits through phosphorylation of PJA2 which binds and ubiquitinates these subunits, leading to their subsequent proteolysis.

#### **Immunogen information**

Immunogen: protein kinase, cAMP-dependent, catalytic, beta

Synonyms: None

Observed MW: 46-55 kDa UniprotID: P22694

#### **Application**

Reactivity: Human, Mouse, Rat

## 1

#### Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)


Tel:(0086)027-87384275 Fax: (0086)027-87800889 <u>www.fn-test.com</u>




Tested Application: ELISA, WB, IHC, IF, IP

Recommended dilution: WB: 1:500-1:2000; IP: 1:200-1:1000; IHC: 1:20-1:200; IF: 1:20-1:200

Image:




Immunohistochemistry of paraffin-embedded human medulloblastoma using FNab06778(PRKACB antibody) at dilution of 1:50



IP Result of anti-PRKACB (IP:FNab06778, 4ug; Detection:FNab06778 1:500) with mouse brain tissue lysate 2640ug.





HT-1080 cells were subjected to SDS PAGE followed by western blot with FNab06778(PRKACB antibody) at dilution of 1:1000