

anti- RNF2 antibody

Product Information

Catalog No.:	FNab07348
Size:	100µg
Form:	liquid
Purification:	Immunogen affinity purified
Purity:	≥95% as determined by SDS-PAGE
Host:	Rabbit
Clonality:	polyclonal
Clone ID:	None
IsoType:	IgG
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months (Avoid repeated freeze / thaw cycles.)

Background

Polycombgroup (PcG) of proteins form the multiprotein complexes that are important for the transcription repression of various genes involved in development and cell proliferation. The protein encoded by this gene is one of the PcG proteins. It has been shown to interact with, and suppress the activity of, transcription factor CP2 (TFCP2/CP2). Studies of the mouse counterpart suggested the involvement of this gene in the specification of anterior-posterior axis, as well as in cell proliferation in early development. This protein was also found to interact with huntingtin interacting protein 2 (HIP2), an ubiquitin-conjugating enzyme, and possess ubiquitin ligase activity.

Immunogen information

Immunogen:	ring finger protein 2
Synonyms:	BAP1, DING, HIPI3, RING1B
Observed MW:	39 kDa
UniprotID :	Q99496

Application

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve. East Lake High-Tech Development Zone. Wuhan, Hubei, China(430206)

Tel : (0086)027-87384275


Fax: (0086)027-87800889 www.fn-test.com

Reactivity: Human, Mouse, Rat


Tested Application: ELISA, WB, IHC, IP

Recommended dilution: WB: 1:500 - 1:2000; IHC: 1:50 - 1:200; IP: 1:50 - 1:200


Image:



Immunohistochemistry of paraffin-embedded human prostate cancer using FNab07348(RNF2 Antibody) at dilution of 1:100



IP Result of anti-RNF2 (IP:FNab07348, 4ug; Detection:FNab07348 1:800) with K-562 cells lysate 1200ug.



mouse kidney tissue were subjected to SDS PAGE followed by western blot with FNab07348(RNF2 antibody) at dilution of 1:1000