

## anti- SCML4 antibody

### Product Information

|               |   |
|---------------|---|
| Catalog No.:  | FNab07641   |
| Size:         | 100µg   |
| Form:         | liquid  |
| Purification: | Immunogen affinity purified   |
| Purity:       | ≥95% as determined by SDS-PAGE  |
| Host:         | Rabbit  |
| Clonality:    | polyclonal  |
| Clone ID:     | None  |
| IsoType:      | IgG   |
| Storage:      | PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months (Avoid repeated freeze / thaw cycles.) |

### Background

Putative Polycombgroup(PcG) protein. PcG proteins act by forming multiprotein complexes, which are required to maintain the transcriptionally repressive state of homeotic genes throughout development(By similarity).

### Immunogen information

|              |                                       |
|--------------|---------------------------------------|
| Immunogen:   | sex comb on midleg-like 4(Drosophila) |
| Synonyms:    | SCML4                                 |
| Observed MW: | 50-60 kDa                             |
| UniprotID :  | Q8N228                                |

### Application


|                          |              |
|--------------------------|--------------|
| Reactivity:              | Human, Mouse |
| Tested Application:      | ELISA, WB    |
| Recommended dilution:WB: | 1:500-1:2000 |
| Image:                   |              |

### Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)

Tel : (0086)027-87384275

Fax: (0086)027-87800889 [www.fn-test.com](http://www.fn-test.com)



mouse liver tissue were subjected to SDS PAGE followed by western blot with FNab07641( SCML4 Antibody) at dilution of 1:600