

anti- SF3A3 antibody

Product Information

Catalog No.: FNab07776

Size: 100μg Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

Clone ID: None IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

Background

Subunit of the splicing factor SF3A required for 'A' complex assembly formed by the stable binding of U2 snRNP to the branchpointsequence(BPS) in pre-mRNA. Sequence independent binding of SF3A/SF3B complex upstream of the branch site is essential, it may anchor U2 snRNP to the pre-mRNA. May also be involved in the assembly of the 'E' complex.

Immunogen information

Immunogen: splicing factor 3a, subunit 3, 60kDa

Synonyms: SAP61
Observed MW: 59 kDa
UniprotID: Q12874

Application

Reactivity: Human, Mouse, Rat

Tested Application: ELISA, WB, IHC, IF, IP

Recommended dilution: WB: 1:500-1:2000; IP: 1:200-1:1000; IHC: 1:20-1:200; IF: 1:20-1:200

Wuhan Fine Biotech Co., Ltd.


B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)

Tel:(0086)027-87384275 Fax: (0086)027-87800889 <u>www.fn-test.com</u>


1



Image:




Immunohistochemistry of paraffin-embedded human lung cancer using FNab07776(SF3A3 antibody) at dilution of 1:50



IP Result of anti-SF3A3 (IP:FNab07776, 4ug; Detection:FNab07776 1:600) with Raji cells lysate 2000ug.





Raji cells were subjected to SDS PAGE followed by western blot with FNab07776(SF3A3 antibody) at dilution of 1:300

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve. East Lake High-Tech Development Zone. Wuhan, Hubei, China (430206)

Tel: (0086)027-87384275 Fax: (0086)027-87800889 www.fn-test.com