

anti- SKAP2 antibody

Product Information

Catalog No.: FNab07890

Size: 100μg Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

Clone ID: None IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

Background

The protein encoded by this gene shares homology with Src kinase-associated phosphoprotein 1, and is a substrate of Src family kinases. It is an adaptor protein that is thought to play an essential role in the Src signaling pathway, and in regulating proper activation of the immune system. This protein contains an amino terminal coiled-coil domain for self-dimerization, a plecskstrin homology (PH) domain required for interactions with lipids at the membrane, and a Src homology (SH3) domain at the carboxy terminus. Some reports indicate that this protein inhibits actin polymerization through interactions with actin assembly factors, and might negatively regulate the invasiveness of tumors by modulating actin assembly. Alternative splicing results in multiple transcript variants encoding different isoforms.

Immunogen information

Immunogen: src kinase associated phosphoprotein 2
Synonyms: PRAP, RA70, SAPS, SCAP2, SKAP55R

Observed MW: 50 kDa
UniprotID: 075563

Application

1

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)


Tel:(0086)027-87384275 Fax: (0086)027-87800889 www.fn-test.com




Reactivity: Human, Mouse, Rat Tested Application: ELISA, WB, IHC, IF

Recommended dilution: WB: 1:500 - 1:2000; IHC: 1:50 - 1:100; IF: 1:50 - 1:100

Image:



Immunohistochemistry of paraffin-embedded mouse stomach using FNab07890(SKAP2 Antibody) at dilution of 1:50



human liver tissue were subjected to SDS PAGE followed by western blot with FNab07890(SKAP2 antibody) at dilution of 1:1500