

anti- SND1 antibody

Product Information

Catalog No.: FNab08064

Size: 100μg Form: liquid

Purification: Protein A+G purification

Purity: ≥95% as determined by SDS-PAGE

Host: Mouse

Clonality: monoclonal

Clone ID: 0A1
IsoType: IgG1

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

Background

This gene encodes a transcriptional co-activator that interacts with the acidic domain of Epstein-Barr virus nuclear antigen 2 (EBNA 2), a transcriptional activator that is required for B-lymphocyte transformation. Other transcription factors that interact with this protein are signal transducers and activators of transcription, STATs. This protein is also thought to be essential for normal cell growth. A similar protein in mammals and other organisms is a component of the RNA-induced silencing complex (RISC).

Immunogen information

Immunogen: staphylococcal nuclease and tudor domain containing 1

Synonyms: 100 kDa coactivator, EBNA2 coactivator p100, p100, p100 co activator,

SND1, TDRD11

Observed MW: 102 kDa UniprotID: Q7KZF4

Application

Reactivity: Human, Mouse

1

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)


Tel:(0086)027-87384275 Fax: (0086)027-87800889 <u>www.fn-test.com</u>




Tested Application: ELISA, IF, WB, IHC

Recommended dilution: WB: 1:500 - 1:2000; IHC: 1:50 - 1:200; IF: 1:50 - 1:200

Image:



Immunohistochemistry of paraffin-embedded human kidney cancer using FNab08064(SND1 Antibody) at dilution of 1:100



HeLa cells were subjected to SDS PAGE followed by western blot with FNab08064(SND1 Antibody) at dilution of 1:1000

Wuhan Fine Biotech Co., Ltd.