

anti- SPOP antibody

Product Information

Catalog No.:	FNab08189
Size:	100µg
Form:	liquid
Purification:	Immunogen affinity purified
Purity:	≥95% as determined by SDS-PAGE
Host:	Rabbit
Clonality:	polyclonal
Clone ID:	None
IsoType:	IgG
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months (Avoid repeated freeze / thaw cycles.)

Background

Component of a cullin-RING-based BCR(BTB-CUL3-RBX1) E3 ubiquitin-protein ligase complex that mediates the ubiquitination of target proteins, leading most often to their proteasomal degradation. In complex with CUL3, involved in ubiquitination and proteasomal degradation of BRMS1, DAXX, PDX1/IPF1, GLI2 and GLI3. In complex with CUL3, involved in ubiquitination of H2AFY and BMI1; this does not lead to their proteasomal degradation. Inhibits transcriptional activation of PDX1/IPF1 targets, such as insulin, by promoting PDX1/IPF1 degradation. The cullin-RING-based BCR(BTB-CUL3-RBX1) E3 ubiquitin-protein ligase complex containing homodimeric SPOP has higher ubiquitin ligase activity than the complex that contains the heterodimer formed by SPOP and SPOPL.

Immunogen information

Immunogen:	speckle-type POZ protein
Synonyms:	HIB homolog 1, Roadkill homolog 1, speckle type POZ protein, SPOP, TEF2
Observed MW:	42 kDa
UniprotID :	O43791

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve. East Lake High-Tech Development Zone. Wuhan, Hubei, China(430206)

Tel : (0086)027-87384275

Fax: (0086)027-87800889 www.fn-test.com


Application

Reactivity: Human, Mouse, Rat


Tested Application: ELISA, WB, IHC, IP

Recommended dilution: WB: 1:500-1:2000; IHC: 1:50-1:500; IP : 1:500-1:1000

Image:



Immunohistochemistry of paraffin-embedded human prostate cancer tissue slide using FNab08189(SPOP Antibody) at dilution of 1:200




IP Result of anti-SPOP (IP:FNab08189, 3ug; Detection:FNab08189 1:500) with HeLa cell lysate 3000ug.

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve. East Lake High-Tech Development Zone. Wuhan, Hubei, China(430206)

Tel : (0086)027-87384275

Fax: (0086)027-87800889 www.fn-test.com



HeLa cells were subjected to SDS PAGE followed by western blot with FNab08189 (SPOP antibody) at dilution of 1:500