

anti- TFAP2C antibody

Product Information

Catalog No.: FNab08619

Size: 100μg Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

Clone ID: None IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

Background

Sequence-specific DNA-binding protein that interacts with inducible viral and cellular enhancer elements to regulate transcription of selected genes. AP-2 factors bind to the consensus sequence 5'-GCCNNNGGC-3' and activate genes involved in a large spectrum of important biological functions including proper eye, face, body wall, limb and neural tube development. They also suppress a number of genes including MCAM/MUC18, C/EBP alpha and MYC. Involved in the MTA1-mediated epigenetic regulation of ESR1 expression in breast cancer.

Immunogen information

Immunogen: transcription factor AP-2 gamma(activating enhancer binding protein 2

gamma)

Synonyms: None
Observed MW: 50 kDa
UniprotID: Q92754

Application

Reactivity: Human, Mouse, Rat

1

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)


Tel:(0086)027-87384275 Fax: (0086)027-87800889 <u>www.fn-test.com</u>




Tested Application: ELISA, WB, IHC, FC, IP

Recommended dilution: WB: 1:500-1:2000; IP: 1:200-1:1000; IHC: 1:20-1:200

Image:




Immunohistochemistry of paraffin-embedded human breast cancer tissue slide using FNab08619(TFAP2C Antibody) at dilution of 1:400 heat mediated antigen retrieved with Tris-EDTA buffer(pH9).



IP Result of anti-TFAP2C (IP:FNab08619, 4ug; Detection:FNab08619 1:1000) with HeLa cells lysate 2800ug.





HeLa cells were subjected to SDS PAGE followed by western blot with FNab08619(TFAP2C antibody) at dilution of 1:1000