

# anti-TNFAIP8L2 antibody

#### **Product Information**

Catalog No.: FNab08820

Size: 100µg Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

Clone ID: None IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

### **Background**

Acts as a negative regulator of innate and adaptive immunity by maintaining immune homeostasis. Negative regulator of Toll-like receptor and T-cell receptor function. Prevents hyperresponsiveness of the immune system and maintains immune homeostasis. Inhibits JUN/AP1 and NF-kappa-B activation. Promotes Fas-induced apoptosis(By similarity).

### **Immunogen information**

Immunogen: tumor necrosis factor, alpha-induced protein 8-like 2

Synonyms: None

Observed MW: 18-21 kDa UniprotID: Q6P589

#### **Application**

Reactivity: Human, Mouse, Rat

Tested Application: ELISA, WB, IHC, IP, IF

Recommended dilution: WB: 1:500-1:2000; IP: 1:200-1:2000; IHC: 1:20-1:200; IF: 1:20-1:200

#### Wuhan Fine Biotech Co., Ltd.


B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)

Tel:(0086)027-87384275 Fax: (0086)027-87800889 www.fn-test.com


1



## Image:




Immunohistochemistry of paraffin-embedded human lymphoma using FNab08820(TNFAIP8L2 antibody) at dilution of 1:50



IP Result of anti-TIPE2 (IP:FNab08820, 3ug; Detection:FNab08820 1:500) with mouse cerebellum tissue lysate 4000ug.





mouse cerebellum tissue were subjected to SDS PAGE followed by western blot with FNab08820(TNFAIP8L2 antibody) at dilution of 1:500

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve. East Lake High-Tech Development Zone. Wuhan, Hubei, China (430206)

Tel: (0086)027-87384275 Fax: (0086)027-87800889 www.fn-test.com