

anti- TXNDC5 antibody

Product Information

Catalog No.: FNab09127

Size: 100μg Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

Clone ID: None IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

Background

This gene encodes a member of the disulfide isomerase (PDI) family of endoplasmic reticulum (ER) proteins that catalyze protein folding and thiol-disulfide interchange reactions. The encoded protein has an N-terminal endoplasmic reticulum (ER)-signal sequence, three catalytically active thioredoxin domains and a C-terminal ER-retention sequence. Its expression is induced by hypoxia and its role may be to protect hypoxic cells from apoptosis. Alternative splicing results in multiple transcript variants. Read-through transcription also exists between this gene and the neighboring upstream BLOC1S5 gene.

Immunogen information

Immunogen: thioredoxin domain containing 5 (endoplasmic reticulum)

Synonyms: EndoPDI, ER protein 46, ERP46, Hcc 2, PDIA15, Thioredoxin like

protein p46, TLP46, TXNDC5, UNQ364

Observed MW: 50 kDa
UniprotID: Q8NBS9

Application

1

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)


Tel:(0086)027-87384275 Fax: (0086)027-87800889 www.fn-test.com




Reactivity: Human, Mouse, Rat Tested Application: ELISA, WB, IHC, IF

Recommended dilution: WB: 1:500 - 1:2000; IHC: 1:50 - 1:200; IF: 1:50 - 1:100

Image:



Immunohistochemistry of paraffin-embedded human colon cancer using FNab09127(TXNDC5 antibody) at dilution of 1:100



HepG2 cells were subjected to SDS PAGE followed by western blot with FNab09127(TXNDC5 antibody) at dilution of 1:1000

Wuhan Fine Biotech Co., Ltd.