

anti- VEGF antibody

Product Information

Catalog No.:	FNab09391
Size:	100µg
Form:	liquid
Purification:	Immunogen affinity purified
Purity:	≥95% as determined by SDS-PAGE
Host:	Rabbit
Clonality:	polyclonal
Clone ID:	None
IsoType:	IgG
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months (Avoid repeated freeze / thaw cycles.)

Background

VEGFA, also named as VEGF or VPF, belongs to the PDGF/VEGF growth factor family. It is a growth factor active in angiogenesis, vasculogenesis and endothelial cell growth. VEGFA induces endothelial cell proliferation, promotes cell migration, inhibits apoptosis and induces permeabilization of blood vessels. It binds to the FLT1/VEGFR1 and KDR/VEGFR2 receptors, heparan sulfate and heparin. Defects in VEGFA are a cause of susceptibility to microvascular complications of diabetes type 1(MVCD1). VEGFA has 13 isoforms with MW 16kd to 45kd. Some isoforms has homodimer form, such as VEGFA189(38kd), VEGFA110(34kd). VEGFA is a 34kd to 42kd, dimeric, disulfide-bound glycoprotein. VEGF-A exists in at least seven homodimeric isoforms. The monomers consist of 121, 145, 148, 165, 183, 189, or 206 amino acids(PMID:15602010).

Immunogen information

Immunogen:	vascular endothelial growth factor A
Synonyms:	L VEGFA, MVCD1, Vascular permeability factor, VEGF, VEGF A, VEGFA, VPF
Observed MW:	15-27kDa, 34-45kDa
Uniprot ID :	P15692

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 Gaoxin Ave. East Lake High-Tech Development Zone. Wuhan, Hubei, China(430206)

Tel : (0086)027-87384275

Fax: (0086)027-87800889

www.fn-test.com


Application

Reactivity: Human, Mouse, Rat


Tested Application: ELISA, IHC, IF, WB

Recommended dilution: WB: 1:200-1:2000; IF: 1:10-1:100; IHC: 1:50-1:500

Image:



Immunohistochemistry of paraffin-embedded human lung cancer tissue slide using FNab09391(VEGFA Antibody) at dilution of 1:200



Immunofluorescent analysis of HepG2 cells, using VEGFA antibody (FNab09391) at 1:25 dilution and Rhodamine-labeled goat anti-rabbit IgG (red).


Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 Gaoxin Ave. East Lake High-Tech Development Zone. Wuhan, Hubei, China(430206)

Tel : (0086)027-87384275

Fax: (0086)027-87800889

www.fn-test.com



PC-3 cells were subjected to SDS PAGE followed by western blot with FNab09391(VEGFA antibody) at dilution of 1:800