

anti- VLDLR antibody

Product Information

Catalog No.:	FNab09416
Size:	100µg
Form:	liquid
Purification:	Immunogen affinity purified
Purity:	≥95% as determined by SDS-PAGE
Host:	Rabbit
Clonality:	polyclonal
Clone ID:	None
IsoType:	IgG
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months (Avoid repeated freeze / thaw cycles.)

Background

Binds VLDL and transports it into cells by endocytosis. In order to be internalized, the receptor-ligand complexes must first cluster into clathrin-coated pits. Binding to Reelin induces tyrosine phosphorylation of Dab1 and modulation of Tau phosphorylation(By similarity).

Immunogen information

Immunogen:	very low density lipoprotein receptor
Synonyms:	VLDLR, VLDL receptor, VLDL R, FLJ35024, CHRMQ1
Observed MW:	160 kDa
Uniprot ID :	P98155

Application

Reactivity:	Human, Mouse, Rat
Tested Application:	ELISA, WB
Recommended dilution:WB:	1:500-1:5000
Image:	


Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 Gaoxin Ave. East Lake High-Tech Development Zone. Wuhan, Hubei, China(430206)

Tel : (0086)027-87384275

Fax: (0086)027-87800889

www.fn-test.com



MCF7 cells were subjected to SDS PAGE
followed by western blot with
FNab09416(VLDLR antibody) at dilution of 1:300