

anti- WDR32 antibody

Product Information

Catalog No.: FNab09489

Size: 100μg Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

Clone ID: None IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

Background

WDR32, containing seven WD repeats, is also termed DDB1-and CUL4-associated factor 10(DCAF10), which may function as a substrate receptor for CUL4-DDB1 E3 ubiquitin-protein ligase complex(PMID: 16949367). WDR32 is able to interact with DDB1(PMID: 16949367). WDR32 is supposed to be implicated in process of ubiquitin-dependent protein degradation system. Two isoforms are described and produced by alternative splicing. This WDR32 antibody majorly detects WDR32 isoform1 with a theoretical MW of 61 kDa.

Immunogen information

Immunogen: WD repeat domain 32

Synonyms: DCAF10, FLJ23201, WD repeat domain 32, WDR32

Observed MW: 61 kDa
UniprotID: Q5QP82

Application

Reactivity: Human, Mouse, Rat

Tested Application: ELISA, IHC, IP, WB, IF

Wuhan Fine Biotech Co., Ltd.


B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)

Tel:(0086)027-87384275 Fax: (0086)027-87800889 www.fn-test.com


1



Recommended dilution: WB: 1:200-1:2000; IP: 1:200-1:1000; IF: 1:20-1:200; IHC: 1:20-1:200 Image:




Immunohistochemistry of paraffin-embedded human testis using FNab09489(WDR32 antibody) at dilution of 1:100



IP Result of anti-WDR32 (IP:FNab09489, 4ug; Detection:FNab09489 1:300) with HepG2 cells lysate 6000ug.





HeLa cells were subjected to SDS PAGE followed by western blot with FNab09489(WDR32 antibody) at dilution of 1:800