

anti- ZAP70 antibody

Product Information

Catalog No.: FNab09582

Size: 100μg Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

Clone ID: None IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

Background

This gene encodes an enzyme belonging to the protein tyrosine kinase family, and it plays a role in T-cell development and lymphocyte activation. This enzyme, which is phosphorylated on tyrosine residues upon T-cell antigen receptor (TCR) stimulation, functions in the initial step of TCR-mediated signal transduction in combination with the Src family kinases, Lck and Fyn. This enzyme is also essential for thymocyte development. Mutations in this gene cause selective T-cell defect, a severe combined immunodeficiency disease characterized by a selective absence of CD8-positive T-cells. Two transcript variants that encode different isoforms have been found for this gene.

Immunogen information

Immunogen: zeta-chain (TCR) associated protein kinase 70kDa

Synonyms: 70 kDa zeta associated protein, FLJ17670, FLJ17679, SRK, STD, Syk

related tyrosine kinase, Tyrosine protein kinase ZAP 70, TZK, ZAP 70,

ZAP70

Observed MW: 69 kDa UniprotID: P43403

1

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)

Tel:(0086)027-87384275 Fax: (0086)027-87800889 <u>www.fn-test.com</u>




Application


Reactivity: Human, Mouse
Tested Application: ELISA, WB, IHC

Recommended dilution: WB: 1:500 - 1:2000; IHC: 1:50 - 1:200

Image:



Immunohistochemistry of paraffin-embedded human amygdalitis using FNab09582(ZAP70 antibody) at dilution of 1:50



Jurkat cells were subjected to SDS PAGE followed by western blot with FNab09582(ZAP70 antibody) at dilution of 1:1000