

anti- C-FOS antibody

Product Information

Catalog No.:	FNab09803
Size:	100µg
Form:	liquid
Purification:	Immunogen affinity purified
Purity:	≥95% as determined by SDS-PAGE
Host:	Rabbit
Clonality:	polyclonal
Clone ID:	None
IsoType:	IgG
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months (Avoid repeated freeze / thaw cycles.)

Background

c-FOS, also named as Proto-oncogene c-Fos and G0/G1 switch regulatory protein 7, is a 380 amino acid protein, which contains 1 bZIP(basic-leucine zipper) domain and belongs to the bZIP family. FOS is expressed at very low levels in quiescent cells. When cells are stimulated to reenter growth, FOS undergo 2 waves of expression, the first one peaks 7.5 minutes following FBS induction. At this stage, the FOS protein is localized endoplasmic reticulum. The second wave of expression occurs at about 20 minutes after induction and peaks at 1 hour. At this stage, the FOS protein becomes nuclear. The calculated molecular weight of FOS is 40 kDa, but Phosphorylated FOS protein is about 60-65 kDa.

Immunogen information

Immunogen:	Proto oncogene c Fos
Synonyms:	AP 1, C FOS, Cellular oncogene fos, c-FOS, FOS, G0S7, Proto oncogene c Fos
Observed MW:	65 kDa
Uniprot ID :	P01100

Application

1

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 Gaoxin Ave. East Lake High-Tech Development Zone. Wuhan, Hubei, China(430206)

Tel : (0086)027-87384275

Fax: (0086)027-87800889


www.fn-test.com

Reactivity: Human, Mouse, Rat


Tested Application: ELISA, IF, IHC, WB

Recommended dilution: WB: 1:500-1:2000; IHC: 1:200-1:800; IF: 1:50-1:500


Image:



Immunohistochemistry of paraffin-embedded mouse brain tissue slide using FNab09803(c-FOS antibody) at dilution of 1:400



Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using FNab09803 (c-FOS antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated Goat Anti-Rabbit IgG(H+L)



mouse embryo tissue were subjected to SDS
PAGE followed by western blot with
FNab09803(c-FOS antibody) at dilution of 1:1000