

# anti-Visfatin antibody

#### **Product Information**

Catalog No.: FNab09847

Size: 100μg Form: liquid

Purification: protein A+G purified

Purity: ≥95% as determined by SDS-PAGE

Host: Mouse

Clonality: monoclonal

Clone ID: 2G6
IsoType: IgG2b

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

### **Background**

Nicotinamide phosphoribosyltransferase(NAMPT) has two usual synonyms termed Visfatin and PBEF. Its primary role is to catalyze the condensation of nicotinamide with 5-phosphoribosyl-1-pyrophosphate to yield nicotinamide mononucleotide, an intermediate in the biosynthesis of NAD, which is the rate limiting component in the mammalian NAD biosynthesis pathway. NAMPT is localized in cytoplasm and expressed in large amounts in bone marrow, liver tissue, and muscle tissues. NAMPT inhibits neutrophil apoptosis in experimental inflammation and clinical sepsis. NAMPT levels are altered in plasma of patients with type 2 diabetes mellitus(T2DM), and it is now evidenced that NAMPT may plays a role in lipid metabolism.

#### **Immunogen information**

Immunogen: nicotinamide phosphoribosyltransferase

Synonyms: 1110035O14Rik, DKFZp666B131, NAmPRTase, NAMPT, PBEF,

PBEF1, Pre B cell enhancing factor, VF, VISFATIN

Observed MW: 52 kDa UniprotID: P43490

## **Application**

1

#### Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)

Tel:(0086)027-87384275 Fax: (0086)027-87800889 <u>www.fn-test.com</u>




Reactivity: Human, Mouse, Rat


Tested Application: ELISA, WB, IHC, IF

Recommended dilution: WB: 1:1000-1:4000; IHC: 1:50-1:500; IF: 1:50-1:500

Image:




Immunohistochemistry of paraffin-embedded human breast cancer tissue slide using FNab09847( Visfatin antibody ) at dilution of 1:200



Immunofluorescent analysis of ( -20°C Ethanol ) fixed HeLa cells using FNab09847( Visfatin antibody ) at dilution of 1:100 and Alexa Fluor 488-congugated Goat Anti-Mouse IgG(H+L)





rat heart tissue were subjected to SDS PAGE followed by western blot with FNab09847( Visfatin antibody ) at dilution of 1:2000

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)

Tel: (0086)027-87384275 Fax: (0086)027-87800889 <u>www.fn-test.com</u>