

anti- HIF1a antibody

Product Information

Catalog No.:	FNab09940
Size:	100µg
Form:	liquid
Purification:	protein A+G purified
Purity:	≥95% as determined by SDS-PAGE
Host:	Mouse
Clonality:	monoclonal
Clone ID:	5H12
IsoType:	IgG1
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months (Avoid repeated freeze / thaw cycles.)

Background

Under normal oxygen conditions, HIF1a is continuously produced and destroyed, in a process involving hydroxylation, interaction with von Hippel-Lindau (VHL) protein, polyubiquitylation and subsequent proteasomal degradation. Under hypoxic conditions, hydroxylation is impaired and HIF1a is stabilized. HIF1a localizes in cytoplasm in normoxia, but it can translocate into nuclear in response to hypoxia. The calculated molecular weight of HIF1a is 93 kDa, but the modified protein HIF1a is about 110-120kDa (PMID: 11698256, .PMID: 7539918).

Immunogen information

Immunogen:	hypoxia inducible factor 1, alpha subunit(basic helix-loop-helix transcription factor)
Synonyms:	BHLHE78, MOP1, PASD8
Observed MW:	120 kDa
Uniprot ID :	Q16665

Application

Reactivity:	Human
-------------	-------

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 Gaoxin Ave. East Lake High-Tech Development Zone. Wuhan, Hubei, China(430206)

Tel : (0086)027-87384275


Fax: (0086)027-87800889

www.fn-test.com


Tested Application: ELISA, IF, WB

Recommended dilution: WB: 1:1000-1:10000; IF: 1:50-1:200

Image:



Immunofluorescent analysis of (-20 °C Ethanol) fixed Cobalt Chloride treated HeLa cells using FNab09940 (HIF1a antibody) at dilution of 1:50 and Alexa Fluor 488-Conjugated Goat Anti-Mouse IgG(H+L)



Cobalt Chloride treated HeLa cells were subjected to SDS PAGE followed by western blot with FNab09940(HIF1a Antibody) at dilution of 1:5000