

anti- COX2 antibody

Product Information

Catalog No.: FNab09973

Size: 100µg Form: liquid

Purification: protein A+G purified

Purity: ≥95% as determined by SDS-PAGE

Host: Mouse

Clonality: monoclonal

Clone ID: 4H7
IsoType: IgG2a

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

Background

Prostaglandin-endoperoxide synthase (PTGS), also known as cyclooxygenase, is the key enzyme in prostaglandin biosynthesis, and acts both as a dioxygenase and as a peroxidase. There are two isozymes of PTGS: a constitutive PTGS1 and an inducible PTGS2, which differ in their regulation of expression and tissue distribution. This gene encodes the inducible isozyme. It is regulated by specific stimulatory events, suggesting that it is responsible for the prostanoid biosynthesis involved in inflammation and mitogenesis.

Immunogen information

Immunogen: prostaglandin-endoperoxide synthase 2

Synonyms: COX 2, COX2, Cyclooxygenase 2, GRIPGHS, hCox 2, PGG/HS, PGH

synthase 2, PGHS 2, PHS 2, PHS II, Prostaglandin G/H synthase 2,

Prostaglandin H2 synthase 2, PTGS2

Observed MW: 68 kDa Uniprot ID: P35354

Application

1

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 Gaoxin Ave. East Lake High-Tech Development Zone. Wuhan, Hubei, China (430206)

Tel :(0086)027-87384275 Fax: (0086)027-87800889 <u>www.fn-test.com</u>




Reactivity: Human, Mouse

Tested Application: ELISA, WB, IHC

Recommended dilution: WB: 1:500-1:2000; IHC: 1:100-1:400

Image:



Immunohistochemistry of paraffin-embedded human lung cancer tissue slide using FNab09973(COX2 Antibody) at dilution of 1:200

—100kd

—70kd

-55kd

-40kd

—35kd

-25kd

LPS treated Hela cells were subjected to SDS PAGE followed by western blot with FNab09973(COX2 antibody) at dilution of 1:1000