

anti- NCOR2 antibody

Product Information

Catalog No.: FNab10069

Size: 100μg Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

Background

Transcriptional corepressor. Mediates the transcriptional repression activity of some nuclear receptors by promoting chromatin condensation, thus preventing access of the basal transcription. Isoform 1 and isoform 5 have different affinities for different nuclear receptors. Involved in the regulation BCL6-dependent of the germinal center(GC) reactions, mainly through the control of the GC B-cells proliferation and survival.

Immunogen information

Immunogen: nuclear receptor co-repressor 2

Synonyms: CTG repeat protein 26, CTG26, N CoR2, NCOR2, Nuclear receptor

corepressor 2, SMAP270, SMRT, SMRTE, SMRTE tau, T3 receptor

associating factor, TNRC14, TRAC, TRAC 1, TRAC1

Observed MW: 180 kDa, 275 kDa

Uniprot ID: Q9Y618

Application

Reactivity: Human, mouse, Rat Tested Application: ELISA, WB, IHC

1

Wuhan Fine Biotech Co., Ltd.


B9 Bld, High-Tech Medical Devices Park, No. 818 Gaoxin Ave. East Lake High-Tech Development Zone. Wuhan, Hubei, China(430206)

Tel :(0086)027-87384275 Fax: (0086)027-87800889 <u>www.fn-test.com</u>




Recommended dilution: WB: 1:500-1:2000; IHC: 1:50-1:200

Image:



Immunohistochemistry of paraffin-embedded human lung using FNab10069(NCOR2 Antibody) at dilution of 1:100



LO2 cells were subjected to SDS PAGE followed by western blot with FNab10069(NCOR2 antibody) at dilution of 1:1000