

# anti- TRADD antibody

#### **Product Information**

Catalog No.: FNab10260

Size: 100µg Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

Clone ID: None IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

## **Background**

The nuclear form acts as a tumor suppressor by preventing ubiquitination and degradation of isoform p19ARF/ARF of CDKN2A by TRIP12: acts by interacting with TRIP12, leading to disrupt interaction between TRIP12 and isoform p19ARF/ARF of CDKN2A(By similarity). Adapter molecule for TNFRSF1A/TNFR1 that specifically associates with the cytoplasmic domain of activated TNFRSF1A/TNFR1 mediating its interaction with FADD. Overexpression of TRADD leads to two major TNF-induced responses, apoptosis and activation of NF-kappa-B.

### **Immunogen information**

Immunogen: TNFRSF1A-associated via death domain

Synonyms: Hs.89862, TRADD

Observed MW: 34 kDa, 27 kDa

UniprotID: Q15628

### **Application**

Reactivity: Human, Mouse, Rat Tested Application: ELISA, WB, IHC

### Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)


Tel:(0086)027-87384275 Fax: (0086)027-87800889 www.fn-test.com

1




Recommended dilution: WB: 1:200 - 1:2000; IHC: 1:50 - 1:500

Image:



Immunohistochemistry of paraffin-embedded mouse heart tissue slide using FNab10260( TRADD antibody at dilution of 1:200



Jurkat cells were subjected to SDS PAGE followed by western blot with FNab10260( TRADD Antibody) at dilution of 1:1000