

anti-MMP16 antibody

Product Information

Catalog No.: FNab10494

Size: 100μg Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

Background

Proteins of the matrix metalloproteinase (MMP) family are involved in the breakdown of extracellular matrix in normal physiological processes, such as embryonic development, reproduction, and tissue remodeling, as well as in disease processes, such as arthritis and metastasis. Most MMP's are secreted as inactive proproteins which are activated when cleaved by extracellular proteinases. The encoded protein activates MMP2 by cleavage. This gene was once referred to as MT-MMP2, but was renamed as MT-MMP3 or MMP16.

Immunogen information

Immunogen: Matrix metalloproteinase-16

Synonyms: MMP16, C8orf57, MMP-X2, MT-MMP2, MT-MMP3, MT3-MMP

Observed MW: 60 kDa Uniprot ID: P51512

Application

Reactivity: Human, Mouse, Rat Tested Application: ELISA, WB, IHC, IF

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 Gaoxin Ave. East Lake High-Tech Development Zone. Wuhan, Hubei, China (430206)


Tel :(0086)027-87384275 Fax: (0086)027-87800889 <u>www.fn-test.com</u>

1




Recommended dilution: WB: 1:500-1:2000; IHC: 1:50-1:200; IF: 1:50-1:200

Image:



Immunohistochemistry of paraffin-embedded human breast cancer slide using FNab10494(MMP16 Antibody) at dilution of 1:200



HT-1080 cells were subjected to SDS PAGE followed by western blot with FNab10494(MMP16 Antibody) at dilution of 1:1000