

# anti- CD79A antibody

#### **Product Information**

Catalog No.: FNab10737

Size: 100μg Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

Clone ID: None IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

### **Background**

Required in cooperation with CD79B for initiation of the signal transduction cascade activated by binding of antigen to the B-cell antigen receptor complex(BCR) which leads to internalization of the complex, trafficking to late endosomes and antigen presentation. Also required for BCR surface expression and for efficient differentiation of pro-and pre-B-cells. Stimulates SYK autophosphorylation and activation. Binds to BLNK, bringing BLNK into proximity with SYK and allowing SYK to phosphorylate BLNK. Also interacts with and increases activity of some Src-family tyrosine kinases. Represses BCR signaling during development of immature B-cells.

#### **Immunogen information**

Immunogen: CD79a molecule, immunoglobulin-associated alpha

Synonyms: CD79A, Ig alpha, IGA, MB 1, MB 1 membrane glycoprotein, MB1,

Surface IgM associated protein

Observed MW: 40 kDa Uniprot ID: P11912

## **Application**

1

#### Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 Gaoxin Ave. East Lake High-Tech Development Zone. Wuhan, Hubei, China(430206)


Tel:(0086)027-87384275 Fax: (0086)027-87800889 <u>www.fn-test.com</u>




Reactivity: Human, Mouse, Rat
Tested Application: ELISA, WB, IHC

Recommended dilution: WB: 1:500-1:2000; IHC: 1:20-1:200

Image:



Immunohistochemistry of paraffin-embedded rat spleen tissue slide using FNab10737(CD79A antibody) at dilution of 1:100



Mouse spleen were subjected to SDS PAGE followed by western blot with FNab10737(CD79A antibody) at dilution of 1:1000