

anti- NEIL1 antibody

Product Information

Catalog No.: FNab10789

Size: 100μg Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

Background

Involved in base excision repair of DNA damaged by oxidation or by mutagenic agents. Acts as DNA glycosylase that recognizes and removes damaged bases. Has a preference for oxidized pyrimidines, such as thymine glycol, formamidopyrimidine(Fapy) and 5-hydroxyuracil. Has marginal activity towards 8-oxoguanine. Has AP(apurinic/apyrimidinic) lyase activity and introduces nicks in the DNA strand. Cleaves the DNA backbone by beta-delta elimination to generate a single-strand break at the site of the removed base with both 3'-and 5'-phosphates. Has DNA glycosylase/lyase activity towards mismatched uracil and thymine, in particular in U:C and T:C mismatches. Specifically binds 5-hydroxymethylcytosine(5hmC), suggesting that it acts as a specific reader of 5hmC.

Immunogen information

Immunogen: nei endonuclease VIII-like 1(E. coli)

Synonyms: DNA glycosylase/AP lyase Neil1, Endonuclease 8 like 1, Endonuclease

VIII like 1, FPG1, hFPG1, NEH1, Nei homolog 1, Nei like protein 1,

NEI1, NEIL1

Observed MW: 44kd Uniprot ID: Q96FI4

1

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 Gaoxin Ave. East Lake High-Tech Development Zone. Wuhan, Hubei, China (430206)

Tel:(0086)027-87384275 Fax: (0086)027-87800889 <u>www.fn-test.com</u>




Application

Reactivity: Human, Rat
Tested Application: ELISA, WB

Recommended dilution: WB: 1:500-1:1000

Image:



rat liver were subjected to SDS PAGE followed by western blot with FNab10789(NEIL1 antibody) at dilution of 1:500